

PERTUSSIS

Basis of diagnosis

Clinical criteria	
Fits clinical description	Select the 'Yes' option if the case fits the clinical description as follows: A clinically compatible case characterised by a cough, and one or more of: <ul style="list-style-type: none"> • paroxysms of cough • cough ending in vomiting, cyanosis or apnoea • inspiratory whoop.
Clinical features	Ideally, obtain information on all of the clinical features listed. If the feature was present, record by selecting the 'Yes' option. If not, Select the 'No' option. If not known or unavailable, then Select the 'Unknown' box.
Laboratory criteria	
Laboratory results	Indicate the status of each laboratory testing method. If the laboratory test results were positive select the 'Yes' option, if negative select the 'No' option. If the results of the laboratory test are not yet available, select 'Awaiting results'. If any of the laboratory tests were not carried out, select 'Not Done'.
Epidemiological criteria	
Contact with a confirmed case	Indicate whether the person has had any contact with a confirmed case of pertussis. If not known or unavailable, then select the 'Unknown' option.

Classification

Classification	<p>Under investigation - a case which has been notified but information is not yet available to classify it as probable or confirmed.</p> <p>Probable –a clinically compatible illness AND</p> <ul style="list-style-type: none"> • a cough lasting 14 days or more. OR • exposure as part of an outbreak (e.g. an institutional outbreak or community-wide outbreak [when there is limited access to testing]). <p>Confirmed – a case that has laboratory definitive evidence OR a clinically compatible illness that is epidemiologically linked to a confirmed case.</p> <p>Laboratory definitive evidence for a confirmed case requires: detection of <i>Bordetella pertussis</i> nucleic acid by PCR or isolation of <i>Bordetella pertussis</i>.</p> <p>Not a case – a case that has been investigated, and subsequently found not to meet the case definition.</p>
----------------	--

Risk factors

Attendance at school, pre-school or childcare	Indicate whether the case attends school, pre-school or childcare. If not known or unavailable, then select the 'Unknown' option.
Other risk factor for pertussis	Specify any other risk factors under surveillance for pertussis if they were present.

Protective factors

Was case immunised	Indicate whether the patient had been immunised with a pertussis-containing vaccine at any time before becoming ill .
Vaccine details	If the case has been immunised (prior to becoming ill), record the details of each dose they received. Record the date of each dose or the age when it was given. Specify the age units (weeks, months, years) by selecting one of the boxes. If the case only received one dose, record the details for the dose received and select the 'Not given' option for the second dose to indicate no further doses had been received. If vaccination dose information is not known or unavailable, then select the 'Unknown' option. Indicate whether the source of immunisation information was patient/caregiver recall or documented for each dose.
Maternal vaccination	If the case is aged less than 5 years, indicate whether the birthing parent was given a pertussis vaccine during pregnancy. If yes, provide the date of vaccination.

Management

Case management	
Exclusion from work or school/pre-school/childcare	Indicate whether the case was excluded from work or school/pre-school/childcare for the appropriate period (refer to the Communicable Disease Control Manual). If the case does not attend work or school/pre-school/childcare select the 'NA' option. If not known or unavailable, then select the 'Unknown' option.
Contact management	
Contacts aged under 7 years	Indicate whether contacts under 7 years of age who were not fully immunised were encouraged to be immunised. If the case had no contacts under 7 years, select the 'NA' (not applicable) option. If not known or unavailable, then select the 'Unknown' option.
Household contacts aged <1 year	Indicate whether the case had any household less than 1 year old. If not known or unavailable, then select the 'Unknown' option. Record the number of household contacts under 1 year, how many of them have had pertussis and how many were offered erythromycin. Note: the number who have had pertussis and/or were given erythromycin should be a subset of the number of household contacts under 1 year.